GPPSS COVID-19 Screening Tool for Families



Parents and guardians, use this checklist every day before sending your children to school.

Does your child have any new, unusual, or worsening symptoms as listed below?

- · Fever or chills
- Cough
- · Shortness of breath or difficulty breathing
- Fatique
- · Muscle or body aches
- Headache

- New loss of taste or smell
- Sore throat
- · Congestion or runny nose
- · Nausea or vomiting
- Diarrhea

Does your child have at least 2 of these symptoms?

- ☐ YES ☐ NO
- ▶ If the answer to the question above is "yes", keep your child home and consult your primary care physician. If a doctor determines that the symptoms are due to another diagnosis, or COVID-19 is ruled out, your child may return to school after being fever-free for 24 hours without the use of fever-reducing medications.
- ► If your child tests positive for COVID-19 OR if your child is exposed to someone who has tested positive for COVID-19, please use the chart below for guidance on isolation and quarantine. Be sure to reach out to your student's school office to report a positive case of COVID-19.

MEDHHS Who is Impacted **Public Health Recommendations** Isolate at home for 5 days (day "0" is day Any individual who tests symptoms begin or day test was taken for positive for COVID-19 and/or those without symptoms); and displays COVID-19 symptoms COVID-19 (without an alternate If symptoms have improved or no symptoms diagnosis or negative Infection developed, may leave isolation after day 5 COVID-19 test) regardless of and wear a well-fitting mask, for 5 more days vaccination status. (ending after day 10).* Wear a well-fitting Monitor symptoms mask around others for for 10 days. 10 days after exposure. Close contact exposed COVID-19 to someone with COVID-19, regardless of Avoid unmasked activities Test 5 days after **Exposure** vaccination status. or activities with higher exposure and if risk of exposing symptoms develop. vulnerable individuals.** * You may remove your mask sooner than day 10 with two sequential negative antigen tests 48 hours apart. If a mask cannot be worn, 10 days of home isolation is recommended.