2016 AP® CHEMISTRY FREE-RESPONSE QUESTIONS

 $Ba^{2+}(aq) + EDTA^{4-}(aq) \rightleftharpoons Ba(EDTA)^{2-}(aq) \qquad K = 7.7 \times 10^7$

- 6. The polyatomic ion $C_{10}H_{12}N_2O_8^{4-}$ is commonly abbreviated as EDTA⁴⁻. The ion can form complexes with metal ions in aqueous solutions. A complex of EDTA⁴⁻ with Ba²⁺ ion forms according to the equation above. A 50.0 mL volume of a solution that has an EDTA⁴⁻(*aq*) concentration of 0.30 *M* is mixed with 50.0 mL of 0.20 *M* Ba(NO₃)₂ to produce 100.0 mL of solution.
 - (a) Considering the value of K for the reaction, determine the concentration of $Ba(EDTA)^{2-}(aq)$ in the 100.0 mL of solution. Justify your answer.
 - (b) The solution is diluted with distilled water to a total volume of 1.00 L. After equilibrium has been reestablished, is the number of moles of $Ba^{2+}(aq)$ present in the solution greater than, less than, or equal to the number of moles of $Ba^{2+}(aq)$ present in the original solution before it was diluted? Justify your answer.

© 2016 The College Board. Visit the College Board on the Web: www.collegeboard.org.